Clozapine (Clozaril®) ALAMEDA COUNTY BEHAVIORAL HEALTH CARE SERVICES Client's Name Date **Initial Medication Instructions** Schizophrenia, schizoaffective disorder, bipolar disorder or other What does this medication help to treat? conditions. Symptoms of these conditions are: □ Poor concentration ☐ Hallucinations (hearing voices) Rapid thoughts ☐ Delusions (beliefs that are false) ☐ Pacing and restlessness ☐ Fearful feelings ☐ Fluctuations in mood ☐ Paranoia or suspiciousness ☐ Insomnia ☐ Agitation, aggression, or hostility Clozapine is sometimes prescribed for other uses; ask your health care professional for more information. ☐ Lack of energy or motivation Other information This medication can take up to 12 weeks to achieve desired results but you should begin to see improvement within the first two weeks of treatment. It is very important to keep all appointments with your clinic, prescriber, and laboratory. Your blood will have to be monitored weekly for the first 6 months of therapy with clozapine. **Side Effects and Management** Common (greater than 10 in 100 clients on this medication) Use caution when driving or operating machinery. Ask your prescriber about taking your dose at **Drowsiness** bedtime if drowsiness occurs. Arise slowly from chairs. Dangle feet off the side of the bed before getting up. **Dizziness** Nausea, Vomiting or Upset Stomach Try taking your clozapine with food, if vomiting continues, call your prescriber. You may take aspirin or Tylenol to relieve your headache. The headache side effect of this medication Headache should subside over time. Talk to your prescriber if headaches are severe or persist for more than one Talk to your prescriber about this side effect. The dose of your medication may need to be lowered. Blurred vision Increased appetite, Weight gain Avoid foods high in fat and sugar. Eat balanced meals and maintain an active lifestyle. Individuals may gain 10-15 pounds over 1-2 months. Constipation Drink plenty of water and increase fiber in your diet. Consult your prescriber about other treatment. Rapid heartbeat Treatment is available; if you experience a rapid heartbeat, consult your prescriber. This may occur in the nighttime hours. Treatment is available if you feel this is bothersome. Consult Increased watering of the mouth with your prescriber. **Uncommon (~1 to 10 in 100 clients on this medication)** Blood abnormality Tell your prescriber if you start to have many infections, or you feel flu-like symptoms, this may be a rare side effect of clozapine in which your immune system may be lowered. To maintain safety, the white blood cell count for all patients taking clozapine must be checked each week for the first six months and then every other week after that. Alert your prescriber immediately. You may be at greater risk if you are taking higher doses or if Seizures vou take other drugs or substances that can cause seizures. Increased sensitivity to the sun Avoid unnecessary or prolonged exposure to sunlight and wear protective clothing, sunglasses, and sunscreen. Muscle Side Effects, You may experience muscle stiffness, spasm, or restlessness. If this should happen, tell your Restlessness prescriber so that your dose may be adjusted or a medication can be given to counteract these Rare (less than 1 in 100 clients on this medication) Call your prescriber immediately if you experience any of these: Seizures or convulsions

- Fine worm-like tongue movements, unusual face, mouth, or iaw movements
- Fast or irregular heartbeat (palpitations)
- Slow or difficult speech
 - **References: Clinical Psychopharmacology, LexiComp 2011**
- Shuffling walk
- Blurred vision
- Fever
- Skin rash or vellowing of the skin or eyes
- Difficulty swallowing or breathing
 - Risk of stroke (in elderly individuals)

Note: This information is not intended to cover all possible uses, precautions, interactions or adverse effects for this medication. If you have further questions about the medications you are taking check with your health care professional.